

AD-A261 079

OFFICE OF NAVAL RESEARCH Grant No. N00014-90-J-1263 RCT Project 4133002---05 Technical Report #25

A NUMERICAL SIMULATIONS OF ADSORPTION ONTO A CRYSTALLINE SURFACE

by

J. A. Hernando * and L. Blum**

Prepared for Publication in the J. Phys. C.



*Comisión Nacional de Energía Atómica, Buenos Aires, Argentina

and



**Department of Physics, University of Puerto Rico P.O. Box 23343 Río Piedras, P.R. 00931-3343, USA

Reproduction in whole or in part is permitted for any purpose of the United States Government

*This document has been approved for public release and sale; its distribution is unlimited

*This statement should also appear in Item 10 of Document Control Data - DD Form 1473. Copies of form available from congnizant contract administrator.

| SECURITY CONSINCATION OF THIS PAGE | | | | | | |
|--|--|--|--|------------------|--------------|--|
| REPORT DOCUMENTATION PAGE | | | | | | |
| 1a. REPORT SECURITY CLASSIFICATION | | 16 RESTRICTIVE MARKINGS | | | | |
| 2a. SECURITY CLASSIFICATION AUTHORITY | | 3 DISTRIBUTION AVAILABILITY OF REPORT | | | | |
| 2b. DECLASSIFICATION / DOWNGRADING SCHEDULE | | | | | | |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S) | | 5 MONITORING ORGANIZATION REPORT NUMBER(S) | | | | |
| Technical Report #25 | | | | | | |
| Physics Department 6b. OFFICE SYMBO (If applicable) | | 78 NAME OF MONITORING ORGANIZATION | | | | |
| University of Puerto Rico | | | | | | |
| 6c. ADDRESS (City, State, and ZIP Code) | 7b ADDRESS (City, State, and ZIP Code) | | | | | |
| Río Piedras Puerto Rico 00931-3343 | | | | | | |
| 8a. NAME OF FUNDING SPONSORING ORGANIZATION Chemistry | · · · · · · · · · · · · · · · | | 9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER | | | |
| ORGANIZATION Chemistry Office of Naval Research | Code 472 | RCT Proje | ct 4133002 | -05 | | |
| 8c. ADDRESS (City, State, and ZIP Code) | | 10 SOURCE OF FUNDING NUMBERS | | | | |
| Arlington | | PROGRAM | PROJECT | TASK | WORK UNIT | |
| Virginia 22217-5000 | | ELEMENT NO | NO. | NO | ACCESSION NO | |
| | <u> </u> | <u> </u> | <u> </u> | <u> </u> | | |
| 11. TITLE (Include Security Classification) A NUMERICAL SIMULATION OF ADSORPTION ONTO A CRYSTALLINE SURFACE | | | | | | |
| 12. PERSONAL AUTHOR(S) J.A. Hernando and L. Blum | | | | | | |
| 13a. TYPE OF REPORT 13b. TIME COVERED FROM TO | | 14. DATE OF REPORT (Year, Month, Day) 15. PAGE COUNT 1-15-93 | | | | |
| 16. SUPPLEMENTARY NOTATION | | | | | | |
| | | | | | | |
| 17 COSATI CODES | Continue on reverse | if necessary and | identify by bloc | k number) | | |
| FIELD GROUP SUB-GROUP | 4 | | | | | |
| | 4 | | | | | |
| 19 ARSTRACT (Continue on reverse if necessary | and identify by block i | number) | | | | |
| 19 ABSTRACT (Continue on reverse if necessary and identify by block number) | | | | | | |
| A particular model for the adsorption of a dense fluid onto a crystalline surface has been studied (L. Blum | | | | | | |
| and D.A. Huckaby, J. Chem. Phys 94, 6887 (1991). The fluid is arbitrary and the crystalline surface is | | | | | | |
| modelled by a hard wall decorated with a triangular lattice of sticky sites. This model can be mapped on a | | | | | | |
| lattice gas model with interaction potentials related to the particle distribution functions of the smooth (undecorated) wall problem. Approximate adsorption isotherms in the 8-p plane (fraction of occupied sites | | | | | | |
| vs. bulk density) have also been drawn. However, the effect that the three particle distribution function has | | | | | | |
| on the critical point has also been estimated (D.A. Huckaby and L. Blum, submitted) and it is quite important. | | | | | | |
| Here we have performed Monte Carlo simulations of a model system inspired on this work. It is a 2D systems | | | | | | |
| interacting via a hard sphere plus a triangular well potential and immersed in an external field described by | | | | | | |
| a triangular lattice of sticky sites. We have studied its phase diagram paying some attention also to its | | | | | | |
| behavior in the limit of infinitesimally small sticky adhesion. | | | | | | |
| İ | • | | | | | |
| 20. DISTRIBUTION / AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED SAME AS RPT DTIC USERS 21 ABSTRACT SECURITY CLASSIFICATION | | | | | | |
| UNCLASSIFIED/UNLIMITED SAME AS | 22b TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL | | | | | |
| 228 NAME OF RESPONSIBLE INDIVIDUAL | | | (203) 696–4410 ONR 472 | | | |
| Dr. Robert Nowak | iPR edition may be used u | | | | OF THE 33.55 | |
| DD FORM 1473, 84 MAR 83 A | All other editions are a | | SECURITY | CLASSIFICATION I | OF THIS PAGE | |

A Numerical Simulation of Adsorption onto a Crystalline Surface

J.A.Hernando^a and L.Blum^b

^a Depto de Física, Comisión Nacional de Energía Atómica, Av. Libertador 8250, 1429 Buenos Aires, Argentina

Dept. of Physics, University of Puerto Rico, Rio Piedras, PR 00931-3343

The sticky site model (SSM) has been recently introduced as a simplified but not too unrealistic model of a structured solid-liquid interface [1]. Here we focus our attention on the adsorbed layer onto a surface that is modelled as a smooth plane surface with sticky points on it. These sticky points form a crystalline lattice representing the adsorption sites and the adsorbed layer feels them as an external potential. This potential u(r) can be represented by its Boltzmann factor

$$exp(-\beta u(r)) = 1 + \sum_{n_1,n_2} \lambda \delta(r - n_1 a_1 - n_2 a_2)$$
 (1)

Here a_1,a_2 describe the lattice, the stickiness parameter λ represents the fugacity of an adsorbed atom and $\beta=1/kT$.

We have performed MC simulations on a system of 196 particles that interact between themselves with the hard sphere-triangular well potential

$$v(r) = \begin{cases} \infty & r \leq 1 \\ V_1 \frac{r-1}{r_0-1} & 1 < r \leq r_0 \\ 0 & r > r_0 \end{cases}$$

the sticky sites form a triangular lattice of constant $a = r_0$ and are modelled by the potential

$$\mathbf{u}(\mathbf{r}) = \left\{ \begin{array}{cc} U_{\bullet} & \mathbf{r} \leq \delta \\ 0 & \mathbf{r} > \delta \end{array} \right.$$

with $r_0 = 1.30$, $\delta = 0.15$. There also are 196 sticky sites, the density is $\eta = \pi N/(4V) = .53$ (close packing is et a = .9) and the parameters that vary are T,V_s and U_s. Several data sets were considered and the runs' length was typically between 20 and 30 10° MC steps. In Fig. 1a,b,c,d, we show results for U_s = V_t = -50, T = 15,25,40 and in fig. 1c,f we plot adsorption isotherms as a function of u and adsorption as a function of T with U_s labelling the curves; V_t = -50 always. It is seen that the system undergoes a first order phase transition liquid-solid, that the solid commensurates with the lattice of sticky sites and that the uprising in adsorption coincides with $g_2(r)$ developing shoulders corresponding to the triangular lattice peaks.

References

[1] L. Blum, Adv. Chem. Phys. Vol LXXVIII, edited by I. Prigogine and S.A. Rice, John Wiley and Sons, N. York (1990).

Figure Captions

Figure 1. a) Stuck particles probability distribution function for $U_s=V_t=-50$, T=15,25,40 (solid, dot and dashed lines); b) pair distribution function for T=15 and triangular lattice peak positions (solid and dashed); c) idem as b) with T=25; d) idem as b) with T=40; e) adsorption isotherm as a function of U_s for T=15,25,40; f) adsorption as a function of T for $U_s=-70,-50,-30$.

3

DTIC QUALITY INSPECTED 3

| | | _ | | |
|---------------|-----------|-------------|--|--|
| Accession For | | | | |
| NTIS | GRALI | 3 | | |
| DTIC ' | | | | |
| Unannounced 🔲 | | | | |
| Justification | | | | |
| | | | | |
| Ву | | | | |
| Distribution/ | | | | |
| Avai | lability | Codes | | |
| | Avail and | d/or | | |
| Dist | Special | | | |
| | 1 1 | | | |
| 10~1 | 1 ! | | | |
| 7 | 1 1 | | | |
| 11, | | | | |